

# **LEGIBILITY NOTICE**

A major purpose of the Technical Information Center is to provide the broadest dissemination possible of information contained in DOE's Research and Development Reports to business, industry, the academic community, and federal, state and local governments.

Although a small portion of this report is not reproducible, it is being made available to expedite the availability of information on the research discussed herein.

CONFIDENTIAL  
Received by OSTI

AUG 04 1988

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

LA-UR--88-2023

DE88 014426

TITLE ELASTIC NEUTRON SCATTERING IN  $UCd_{11}$

AUTHOR(S) J. D. Thompson, A. C. Lawson, M. W. McElfresh, A. P. Sattelberger,  
and Z. Fisk

SUBMITTED TO 6th International Conference on Crystal-Field Effects and  
Heavy-Fermion Physics, Frankfurt, West Germany, July 18, 1988,  
~~Los Alamos National Laboratory, Los Alamos, New Mexico 87545~~

### DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. References herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution or to allow others to do so, for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

 **Los Alamos** Los Alamos National Laboratory  
Los Alamos, New Mexico 87545

# ELASTIC NEUTRON SCATTERING IN $\text{UCd}_{11}$

J. D. Thompson, A. C. Lawson, M. W. McElfresh,

A. P. Sattelberger, and Z. Fisk

Los Alamos National Laboratory

Los Alamos, NM 87545 USA

## Abstract

We report results of neutron powder diffraction experiments on the heavy-electron magnet  $\text{UCd}_{11}$  at temperatures above and below its ordering temperature. These measurements set an upper limit on the ordered moment of  $1.5\mu_B/\text{U}$ .

Correspondence to: J. D. Thompson  
Los Alamos National Laboratory  
MS K764  
Los Alamos, NM 87545 USA  
(505-667-6416)

Specific heat measurements on  $\text{UCd}_{11}$  show a large electronic contribution ( $\gamma \simeq 840 \text{ mJ/mole K}^2$ ) above 5K, at which temperature occurs an anomaly indicative of a magnetic phase transition in the bulk [1]. Associated with this anomaly is a very weak feature in the magnetic susceptibility. More recent muon-spin rotation/relaxation measurements [2] suggest antiferromagnetic order at 5K but neither the ordered moment nor magnetic structure could be determined from these experiments. For the two other U-based heavy-electron magnets  $\text{U}_2\text{Zr}_7$  [3] and  $\text{UCu}_5$  [4], neutron scattering experiments have established that the magnetic order is simple antiferromagnetic, with ordered moments of  $0.8 \pm 0.1$  and  $0.9 \pm 0.1 \mu_B/\text{U}$  respectively, values close to 1/3 of their corresponding paramagnetic moments. Interestingly, the factor 1/3 also appears in the ratio of their electronic specific heat  $\gamma$  below  $T_N$  to that above  $T_N$  [5].  $\text{UCd}_{11}$  also exhibits the ratio  $\gamma(T=0)/\gamma(T \gtrsim T_N) = 0.3$  [1] from which, if the analogy holds, we would imply an ordered moment of  $\gtrsim 1 \mu_B/\text{U}$ .

To minimize the problem associated with the extremely large neutron absorption cross-section of naturally abundant Cd, we have performed neutron diffraction on samples enriched to 98.55%  $\text{Cd}^{114}$ , which has a much smaller absorption cross section. An approximately 5 g sample was prepared by sealing under vacuum fine U powder in a quartz tube with distilled  $\text{Cd}^{114}$  and heating to  $470^\circ\text{C}$  for 7 days. X-ray analysis of the reacted powder showed the material to be  $\gtrsim 97\%$   $\text{UCd}_{11}$ , having the  $\text{Pn}3\text{m}$  crystal structure with a lattice parameter  $a_0 = 9.290 \text{ \AA}$ . Magnetic susceptibility measured on 85 mg of the sample showed Curie-Weiss behavior for  $20 < T < 350\text{K}$ , with  $\mu_{\text{eff}} = 3.86 \mu_B/\text{U}$  and  $\theta = -46.6 \text{ K}$ , and a plateau near 5 K, characteristic of the magnetic transition in  $\text{UCd}_{11}$ .

Neutron scattering was performed on the General Purpose Powder Diffractometer at Argonne's Intense Pulsed Neutron Source. For these experiments, the 5 g sample was encapsulated in a thin-walled aluminum holder that was mounted on the cold finger of a  $^4\text{He}$  cryostat whose temperature could be controlled down to 2K. Parts of the sample holder exposed to the neutron beam and not containing sample were masked by a boron-nitride shield. Approximately 20 hours of data were collected at three temperatures 8, 4, and 2K. Each data set was analyzed by Rietveld refinement using the codes developed by Larson and Von Dreele [6].

Results of the measurements at 2K are shown in Fig. 1. In fitting these data, it was necessary to include scattering from the Al sample holder, BN mask, and "displaced" Al arising from scattering off the Al tail of the cryostat. In addition, to the majority phase  $\text{UCd}_{11}$ , we also found evidence for a small amount ( $\sim 2\%$ ) of free Cd, in agreement with the x-ray analysis. By taking these scattering sources into account, all peaks (except one with  $d = 1.409 \text{ \AA}$ ) could be indexed and the refined parameters for  $\text{UCd}_{11}$  are given in Table I. The unidentified peak also appeared in the 4 and 8K spectra. Additional structural details of  $\text{UCd}_{11}$  will be given elsewhere [7] but we note here that assuming unity occupation of the U-site gives a stoichiometry of  $\text{UCd}_{9.5}$  and that no structural change could be detected to within  $\Delta a_0 = \pm 0.00008 \text{ \AA}$ . Further, within the resolution of these experiments, no additional lines could be found below 5K that might be expected to appear because of antiferromagnetic order.

A minimum detectable moment was estimated assuming a simple ferromagnetic structure, which would provide an upper limit. We assume that a magnetic peak would be observed if its intensity were greater than the (3,0,0)/(2,1,1) nuclear peak indicated by the arrow in Fig. 1(c).

Because the structure factor of this peak is known, we can calculate a limit of detectability to be  $1.5 \mu_B/U$ . Therefore, we conclude that the ordered moment of "UCd<sub>11</sub>" must be less than  $1.5 \mu_B/U$ , that the magnetic structure is not simple, or possibly both. The first conclusion is implicated on the basis of analogy to other U-based heavy-electron magnets.

Work at Los Alamos was performed under the auspices of the USDOE. This work also benefited from the use of the Intense Pulsed Neutron Source at Argonne National Laboratory, which is funded by the U. S. Department of Energy, BES-Materials Science, under contract W-31-109-ENG-38. We thank J. A. O'Rourke for the x-ray measurements.

TABLE I  
Crystal Data for  $\text{UCd}_{11}$  at 2 K

Space Group Pm3m			$a_0 = 9.2480(8) \overset{0}{\text{\AA}}$		
Atomic Positions					
Atom	x	y	z	$u(\overset{0}{\text{\AA}^2})$	Fraction
U1	0	1/2	0	0.0072( 5)	1
Cd1	1/2	1/2	1/2	0.0033(13)	0.81(3)
Cd2	0.1567(2)	0.1567(2)	0.1567(2)	0.0063( 6)	0.84(1)
Cd3	0	0.3441(2)	0.3441(2)	0.0082( 5)	0.87(1)
Cd4	1/2	0.2664(1)	0.2664(1)	0.0078( 4)	0.88(1)


### Figure Caption

Fig. 1. Time-of-flight neutron diffraction spectra for  $\text{UCd}_{11}$  at 2K. Data in panels (a) and (b) were collected with detectors at  $2\theta = 148^\circ$  and those in (c) with  $2\theta = 90^\circ$ . Crosses are data, the solid line is the Rietveld fit and the lower curve in (a) and (b) is the residual. Vertical tick marks indicate d-spacings of Bragg reflections for  $\text{UCd}_{11}$ , Al, Cd, BN, and displaced Al, respectively, from the bottom set upwards.



## References

- [1] Z. Fisk, G. R. Stewart, J. O. Willis, H. R. Ott, and F. Hulliger, Phys. Rev. B 30 (1984) 6360.
- [2] S. Barth, H. R. Ott, F. Hulliger, F. N. Gygi, A. Schenck, and T. M. Rice, Hyperfine Interactions 31 (1986) 403.
- [3] D. E. Cox, G. Shirane, S. M. Shapiro, G. Aeppli, Z. Fisk, J. L. Smith, J. Kjems and H. R. Ott, Phys. Rev. B 33 (1986) 3614.
- [4] A. Murasik, S. Ligenza and A. Zygmunt, Phys. Stat. Sol. (a) 23 (1974) K163.
- [5] H. R. Ott and Z. Fisk, in Handbook on the Physics and Chemistry of the Actinides, ed. A. J. Freeman and G. H. Lander (Elsevier, New York, 1987) p. 85.
- [6] A. C. Larson and R. B. Von Dreele, Generalized Crystal Structure Analysis System, LA-UR-86-748 (1986).
- [7] A. C. Lawson, J. D. Thompson, M. W. McElfresh, A. P. Sattelberger and Z. Fisk (unpublished).



REPRODUCED FROM  
BEST AVAILABLE COPY